



The Patent Office

The Patent Office
Cardiff Road
Newport
Gwent
NP9 1RH

I, the undersigned, being an officer duly authorised in accordance with Section 62(3) of the Patents and Designs Act 1907, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the Patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or the inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.


In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


Signed

Dated

10 AUG 1994



For official use



30 SEP 1993

-40CT '934002B8446

FAT 1 77 UC

25.00

Your reference

136420

9320132.5

Notes

Please type, or write in dark ink using CAPITAL letters. A prescribed fee is payable for a request for grant of a patent. For details, please contact the Patent Office (telephone 071-438 4700).

Rule 16 of the Patents Rules 1990 is the main rule governing the completion and filing of this form.

Ⓜ Do not give trading styles, for example, 'Trading as XYZ company', nationality or former names, for example, 'formerly (known as) ABC Ltd' as these are not required.

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977 and will inform the applicant if such prohibition or restriction is necessary. Applicants resident in the United Kingdom are also reminded that under Section 23, applications may not be filed abroad without written permission unless an application has been filed not less than 6 weeks previously in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction revoked.

**The
Patent
Office**

Request for grant of a Patent

Form 1/77

Patents Act 1977

1 Title of invention

- 1 Please give the title of the invention

SYNTHESIS OF 17-(3-PYRIDYL) STEROIDS

2 Applicant's details

☐ First or only applicant

- 2a If you are applying as a corporate body please give:

Corporate name BRITISH TECHNOLOGY GROUP LTD

Country (and State of incorporation, if appropriate) U.K.

- 2b If you are applying as an individual or one of a partnership please give in full:

Surname

Forenames

- 2c In all cases, please give the following details:

Address 101 NEWINGTON CAUSEWAY
LONDON

UK postcode (if applicable) SE1 6BU

Country U.K.

ADP number (if known) 414001

2d, 2e and 2f: If there are further applicants please provide details on a separate sheet of paper.

Second applicant (if any)

2d If you are applying as a corporate body please give:

Corporate name

Country (and State
of incorporation, if
appropriate)

2e If you are applying as an individual or one of a partnership please give full:

Surname

Forenames

2f **In all cases**, please give the following details:

Address

UK postcode
(if applicable)

Country

ADP number
(if known)

Ⓢ An address for service in the United Kingdom must be supplied

Please mark correct box

Ⓢ **Address for service details**

3a Have you appointed an agent to deal with your application?

Yes ☒ No ☐ → go to 3b

↓
please give details below

Agent's name MR. R.K. PERCY, M.A. C.P.A.,

Agent's address BRITISH TECHNOLOGY GROUP LTD
101 NEWINGTON CAUSEWAY
LONDON

Postcode SE1 6BU

Agent's ADP
number

12 40835 07003n.

3b: If you have appointed an agent, all correspondence concerning your application will be sent to the agent's United Kingdom address.

3b If you have not appointed an agent please give a name and address in the United Kingdom to which all correspondence will be sent:

Name

Address

Postcode

ADP number
(if known)

Daytime telephone
number (if available)

- 7 The answer must be 'No' if:
- any applicant is not an inventor
 - there is an inventor who is not an applicant, or
 - any applicant is a corporate body.

8 Please supply duplicates of claim(s), abstract, description and drawing(s).

Please mark correct box(es)

- 9 You or your appointed agent (see Rule 90 of the Patents Rules 1990) must sign this request.

Please sign here ➡

A completed fee sheet should preferably accompany the fee.

7 Inventorship

7 Are you (the applicant or applicants) the sole inventor or the joint inventors?

Please mark correct box

Yes ☐

No ☒

A Statement of Inventorship on Patents Form 7/77 will need to be filed (see Rule 15).

8 Checklist

8a Please fill in the number of sheets for each of the following types of document contained in this application.

Continuation sheets for this Patents Form 1/77

Claim(s)

3

Description

Abstract

1

Drawing(s)

8b Which of the following documents also accompanies the application?

Priority documents (please state how many)

Translation(s) of Priority documents (please state how many)

Patents Form 7/77 – Statement of Inventorship and Right to Grant
(please state how many)

Patents Form 9/77 – Preliminary Examination/Search

Patents Form 10/77 – Request for Substantive Examination

9 Request

I/We request the grant of a patent on the basis of this application.

Signed

R.K. PERCY

Agent for the Applicant

Date

29 SEPTEMBER 1993

(day month year)

Please return the completed form, attachments and duplicates where requested, together with the prescribed fee to either:

☐ The Comptroller
The Patent Office
Cardiff Road
Newport
Gwent
NP9 1RH

or

☐ The Comptroller
The Patent Office
25 Southampton Buildings
London
WC2A 1AY

SYNTHESIS OF 17-(3-PYRIDYL) STEROIDS

Background of the invention

1. Field of the invention


15 This invention relates to the synthesis of steroids which have a 16, 17-double bond and a 17-(3-pyridyl) substituent.

2. Description of the related art

10 In the unpublished patent applications GB 9305269.4 (British Technology Group Ltd.) and PCT GB93/00531 (British Technology Group Ltd., S.E. Barrie, M. Jarman and G.A. Potter), both filed on 15th March 1993, we have described 16,17-ene-17-(3-pyridyl) steroids as a class of compounds useful for treatment of androgen- and oestrogen- dependent disorders, especially prostatic and breast cancer respectively. A few of these
15 compounds have previously been mentioned in the literature as intermediates in synthesis, to prepare final products having other uses, but otherwise these compounds were new. Since that date, posters have been presented, notably at the SmithKline Beecham Research Symposium, Robinson College Cambridge, England, 20 25-26 March 1993 and at the British Association for Cancer Research meeting in Sheffield, England, 28-31 March 1993. The Cambridge poster describes the synthesis of an exemplified 16,17-ene-17-(3-pyridyl) steroid as follows:

25 Synthesis of this molecule was envisaged via a possible palladium catalysed cross-coupling [G.A. Potter and R. McCague, J. Org Chem. 55, 6184-6187 (1990)] of a steroidal enol triflate (trifluoromethylsulfonate) with a suitable 3-pyridyl nucleophilic coupling partner. This was achieved by using diethyl(3-pyridyl)borane in aqueous THF, with sodium carbonate as
30 nucleophilic activator. The reaction proceeded remarkably efficiently, without possible triflate hydrolysis or ethyl coupling, providing the desired compound in 80% isolated yield (Scheme 1):

Scheme 1




(wherein in the final compound R' = H)

15 Triflates are expensive starting materials, so an alternative route is desirable. Further, notwithstanding what might appear from a literal reading of the poster, to obtain the 3-ol the reaction has to be carried out on the 3-acetate as protecting group, not the 3-ol. The 3-acetate is then hydrolysed to the

20 3-ol subsequently in a separate step. Our unpublished patent application proposes an alternative route, as follows (Scheme 2):


Scheme 2



wherein X represents the residue of the A, B and C rings of a steroid, R represents a hydrogen atom or an alkyl group of 1-4 carbon atoms, R¹⁴ represents a hydrogen atom, a halogen atom or an alkyl group of 1 to 4 carbon atoms and each of the R¹⁵ substituents independently represents a hydrogen atom or an alkyl or alkoxy group of 1-4 carbon atoms, a hydroxy group or an alkylcarbonyloxy group of 2 to 5 carbon atoms or together represent an oxo or methylene group or R¹⁴ and one of the R¹⁵ groups together represent a double bond and the other R¹⁵ group represents a hydrogen atom or an alkyl group of 1 to 4 carbon atoms, and R¹⁶ represents a hydrogen atom, halogen atom, or an alkyl group of 1 to 4 carbon atoms, and Z¹ and Z² independently represent hydroxy or alkoxy or alkyl of 1-4 carbon atoms each, preferably ethyl or methoxy.

Summary of the invention

It has now surprisingly been found that in the preparation of the preferred compound, 3 β -acetoxy-17-(3-pyridyl)androsta-5, 16-diene, of formula (1):




(1)

the reaction can be carried out via the vinyl iodide intermediate, but using the unprotected 3 β -hydroxy compound, while keeping the proportion of organoboron compound (borane)

used in the cross-coupling reaction to not more than 1.2 equivalents per equivalent of steroid and that the reaction product can then be isolated without chromatography. This route is therefore amenable to large scale synthesis.

The principle of the invention may be expressed as a method of preparing a 3 β -hydroxy- or 3 β - (lower acyloxy) 16,17-ene-17-(3-pyridyl)-substituted steroid, wherein the 3 β - (lower acyloxy) group of the steroid has from 2 to 4 carbon atoms, which comprises subjecting a 3 β -hydroxy-16,17-ene-17-iodo or -bromo steroid to a palladium complex-catalysed cross-coupling reaction with a (3-pyridyl)-substituted borane, in which the pyridine ring is substituted at the 5-position by an alkyl group of 1 to 4 carbon atoms or is unsubstituted thereat, especially with a said borane of formula (2):



(2)

wherein R is a hydrogen atom or an alkyl group of 1-4 carbon atoms and Z¹ and Z² independently represent hydroxy or alkoxy or alkyl of 1-3 carbon atoms each or Z¹ and Z² together represent an alkylenedioxy group of 2 or 3 carbon atoms, in a proportion of from 1.0 to 1.2 equivalents of boron compound per equivalent of steroid, in an organic liquid, which is a solvent for the 3 β -hydroxy steroidal reaction product, and optionally acylating the 3 β -hydroxy reaction product.

Description of the preferred embodiments

Preferred embodiments of the invention include the features set forth in the claims, q.v.

Preferably the 17-iodo ("vinyl iodide") starting compound has a D-ring with optional substitution in position 14, 15 and/or 16 as shown in Scheme 2. Most preferably it is prepared from the corresponding 17-ketone, conveniently via the corresponding

hydrazone. Preferably the vinyl iodide is unsubstituted in the 14, 15 and 16-positions, in which case it can be prepared from dehydroepiandrosterone (DHEA). In the hydrazination it is preferable to use hydrazine hydrate together with a catalytic amount of a proton provider which is most preferably hydrazine sulfate.

5 The hydrazone is preferably iodinated with iodine or brominated with bromine in the presence of a strong base such as a tetraalkylguanidine, especially tetramethylguanidine which is cheaply and readily available.

10 In the cross-coupling reaction, the boron compound is preferably a diethylborane or a dimethoxyborane ($Z^1=Z^2=Et$ or $-OMe$). Other boranes include those in which the boron atom is part of a cyclic ether ring e.g. as in $Z^1, Z^2 = 1,2$ -ethylenedioxy or 1,3-propylenedioxy. The proportion of borane added is no more than 1.2 moles of boron per mole of steroid, preferably about 1.1. The palladium compound is a palladium (0) phosphine complex such as tetrakis(triphenylphosphine) palladium (0) or a compound reducible to a palladium (0) phosphine species, especially

15 20 bis(triphenylphosphine) palladium (II) chloride.

The cross-coupling reaction is preferably carried out in two phases, one aqueous, one organic. The organic phase comprises an organic solvent for the 3β -hydroxy steroidal reaction product, especially tetrahydrofuran (THF). Other cyclic ethers such as dioxane could be used in place of THF. Preferably, a nucleophilic activator, such as sodium carbonate, is used in which case it is normally present in the aqueous phase. After the reaction, inorganic salts are removed by first adding another organic solvent, preferably diethyl ether, which is a solvent for the organoboron contaminants produced in the reaction product, and miscible with the first-mentioned organic solvent (e.g. THF), but immiscible with water, whereafter the organic, e.g. (THF-diethyl ether), phase and water (aqueous phase) can be separated. After this separation, the THF and diethyl ether are

25 30 35 evaporated as a mixture and the remaining reaction product is


washed with a third organic solvent, which can be diethyl ether, preferably cooled to below room temperature, most especially to 10°C or lower. The 3 β -hydroxy steroid reaction product has a low solubility in the ether, which, importantly, removes the
5 organoboron compound/s (and also the palladium compound/s).

To prepare the 3 β -acyloxy (alkylcarbonyloxy) compounds, standard acylating agents such as acetyl, propionyl or butyryl chloride or anhydride can be used. The method of acylation may require some modification for isolation of the product.

10 The following Example illustrates the invention.

EXAMPLE

Illustrative reaction scheme



(a) Dehydroepiandrosterone-17-hydrazone

To a stirred solution of dehydroepiandrosterone (28.8g, 0.1 mol) in ethanol (500ml) was added hydrazine hydrate (19.5ml, 0.4 mol), followed by a solution of hydrazine sulfate (65mg, 0.5 mmol) in water (2ml). After stirring for 3 days the mixture was poured into water (3 litres) to precipitate the product as a white crystalline solid. The product was collected by filtration on a sinter, washed with cold water (2 x 50ml), then with Et₂O (50ml). The product was then dried in vacuo, firstly over silica gel, and finally over P₂O₅, to give the title compound as a white crystalline solid (29.6g, 98%).

Notes 1) The method of Schweder et al., p. 202, compound No. 2 therein (using triethylamine) gave a very fine crystalline product which was difficult to filter.

2) The method of Schweder et al. p. 203, compound No. 4 therein (using sodium acetate buffer) gave a slightly lower yield (96%) in trial experiments, whereas the modified procedure used above gives a product amenable for filtration, and in excellent yield (98%).

(b) 17-Iodo-androsta-5,16-dien-3 β -ol

To a solution of iodine (53.3g, 0.21 mol) in THF (2l), cooled by an ice/water bath to 0°C was added 1,1,3,3-tetramethylguanidine (63 ml, 57.6g, 0.50 mol).

A solution of dehydroepiandrosterone-17-hydrazone (30.25g, 0.10 mol) in THF (750 ml) was then added slowly to the above iodine solution via a transfer needle over about 2h, whilst maintaining the reaction temperature at 0°C. After all the hydrazone solution was added, the mixture was filtered, and the filtrate concentrated. The remaining oil was then heated on an oil bath for 4h, allowed to cool, and dissolved in Et₂O. The Et₂O solution was washed with 1M HCl until the aqueous phase was acidic, washed with 0.5M NaOH, then 1M Na₂S₂O₃, and finally with water. The Et₂O phase was separated, dried (MgSO₄), and concentrated to give the crude product. Recrystallisation from Et₂O/hexane (3:2) afforded the title compound as off-white crystals (35.8g, 90%).

(c) 17-(3-Pyridyl)androsta-5,16-dien-3 β -ol

Diethyl(3-pyridyl)borane (3.23g, 22 mmol) from Aldrich Chemical Co. Ltd. was added to a stirred solution of 17-iodo-androsta-5,16-dien-3 β -ol (7.96g, 20 mmol) in THF (120ml) containing bis(triphenylphosphine)palladium (II) chloride (140mg, 0.2 mmol). An aqueous solution of sodium carbonate (2M, 50ml) was then added and the mixture heated, with stirring, by an oil bath at 80°C for 48h, and allowed to cool.

The mixture was partitioned between Et₂O and water the organic phase was separated, dried (Na₂CO₃) and twice concentrated from Et₂O by evaporation to remove THF (with Et₂O). The residual solid was then washed with Et₂O (100ml), the Et₂O solution decanted off, and the remaining white solid recrystallised from toluene (3.94g, 56%).

Notes 1) The time required for completion needs to be made longer than when using the vinyl triflate (48h vs 1h) since it has been found that the vinyl iodide reacts much more slowly.

2) It has been found that a smaller excess of borane than described in the earlier application (for the vinyl triflate) aids in isolation of product.

3) The work-up procedure enables the product to be isolated without chromatography, thereby enabling scaling up.

(d) 3 β -Acetoxy-17-(3-pyridyl)androsta-5,16-diene

To a stirred suspension of finely powdered 17-(3-pyridyl)androsta-5,16-dien-3 β -ol (3.50g, 10 mmol) in dry diethyl ether (150ml) containing triethylamine (2.3ml, 16 mmol) and dimethylaminopyridine (0.012g, 0.1 mmol) was added acetyl chloride (1.0ml, 14 mmol). The mixture was then stirred at ambient temperature for 12h, over which time a thick white precipitate of triethylammonium chloride had formed. The mixture was then filtered and the filtrate concentrated to afford the crude product which was recrystallised firstly from ethanol/water (1:1), then finally from hexane to afford the title compound (3.30g, 84%).


Notes 1) This is a modification of standard acetylation conditions [e.g. acetic anhydride or acetyl chloride with pyridine base and dimethylaminopyridine catalyst, see e.g. J. Wicha and M. Masnyk, Bulltein of the Polish Academy of Sciences: Chemistry 33 (1-2), 19-27 (1985)] which uses diethyl ether as solvent with acetyl chloride and triethylamine base so that the by-product triethylammonium chloride, which is insoluble in diethyl ether, is precipitated as it forms in the reaction. The reaction mixture can thus be filtered to remove the precipitated by-product, and evaporation of the filtrate provides the acetylated material.

2) It is essential that the final product, rather than the vinyl iodide or a precursor thereof, is acetylated.

The following claims define some important aspects of the invention, but do not purport to include every conceivable aspect for which protection might be sought in subsequent continuing and foreign patent applications, and should not be construed as detracting from the generality of the inventive concepts hereinbefore described.

CLAIMS

1. A method of preparing a 3 β -hydroxy- or 3 β - (lower acyloxy) 16,17-ene-17-(3-pyridyl)-substituted steroid, wherein the 3 β - (lower acyloxy) group of the steroid has from 2 to 4 carbon atoms, which comprises subjecting a 3 β -hydroxy-16,17-ene-17-iodo or -bromo steroid to a palladium complex-catalysed cross-coupling reaction with a (3-pyridyl)-substituted borane in which the pyridine ring is substituted at the 5-position by an alkyl group of 1 to 4 carbon atoms or is unsubstituted thereat, in a proportion of from 1.0 to 1.2 equivalents of borane per equivalent of steroid, in an organic liquid, which is a solvent for the 3 β -hydroxy steroidal reaction product and, where the 3 β -(lower acyloxy) group is to be prepared, reacting the resulting 3 β -hydroxy steroidal reaction product with an acylating agent effective to replace the hydroxy group by a said lower acyloxy group.
2. A method according to claim 1, wherein the 3 β -hydroxy steroidal reaction product is reacted with an acetylating agent to give the corresponding 3 β -acetoxy-16,17-ene-17-(3-pyridyl) steroid.
3. A method according to claim 1 or 2 wherein the starting steroid has a D-ring of the following partial formula




- wherein Hal is I or Br, X represents the residue of the A, B and C rings of the steroid, R¹⁴ represents the residue of the A, B and C rings of a steroid, R represents a hydrogen atom or an alkyl group of 1-4 carbon atoms, R¹⁴ represents a hydrogen atom, a halogen atom or an alkyl group of 1 to 4 carbon atoms and each of the R¹⁵ substituents independently represents a hydrogen atom or an alkyl or alkoxy group of 1-4 carbon atoms, a hydroxy group

or an alkylcarbonyloxy group of 2 to 5 carbon atoms or together represent an oxo or methylene group or R^{14} and one of the R^{15} groups together represent a double bond and the other R^{15} group represents a hydrogen atom or an alkyl group of 1 to 4 carbon atoms, and R^{16} represents a hydrogen atom, halogen atom, or an alkyl group of 1 to 4 carbon atoms.

4. A method according to claim 3 wherein the starting steroid is 3 β -hydroxyandrost-5-en-17-one dehydroepiandrosterone).

5. A method according to any preceding claim wherein the (3-pyridyl)-substituted borane is of formula:



wherein R is a hydrogen atom or an alkyl group of 1-4 carbon atoms and Z^1 and Z^2 independently represent hydroxy or alkoxy or alkyl of 1-3 carbon atoms each or Z^1 and Z^2 together represent an alkylenedioxy group of 2 or 3 carbon atoms.

6. A method according to any preceding claim wherein the proportion of the borane is about 1.1 equivalent.
7. A method according to any preceding claim in which the reaction is carried out in two phases, one of which is aqueous and the other of which comprises the said organic liquid.
8. A method according to claim 7 wherein the organic liquid is a first organic liquid, the reaction product-containing mixture is worked up by adding a second organic liquid, which is miscible with the first, but immiscible with water, separating the organic and aqueous phases, evaporating the mixture of organic solvents and washing the residue in a third organic liquid under conditions in which the organoboron contaminants are more soluble than the 3β -hydroxy steroidal reaction product.
9. A method according to claim 8 wherein the first organic liquid comprises tetrahydrofuran and the second and third organic liquids comprise diethyl ether.
10. A method according to claim 9 wherein the residue is washed with the third organic liquid cooled to below room temperature.

ABSTRACT

SYNTHESIS OF 17-(3-PYRIDYL) STEROIDS

A method of preparing a 3β -hydroxy- or 3β - (lower acyloxy) 16,17-ene-17-(3-pyridyl)-substituted steroid, wherein the 3β - (lower acyloxy) group of the steroid has from 2 to 4 carbon atoms, which comprises subjecting a 3β -hydroxy-16,17-ene-17-iodo or -bromo steroid to a palladium complex-catalysed cross-coupling reaction with a (3-pyridyl)-substituted borane in which the pyridine ring is substituted at the 5-position by an alkyl group of 1 to 4 carbon atoms or is unsubstituted thereat, in a proportion of from 1.0 to 1.2 equivalents of borane per equivalent of steroid, in an organic liquid, which is a solvent for the 3β -hydroxy steroidal reaction product and, where the 3β - (lower acyloxy) group is to be prepared, reacting the resulting 3β -hydroxy steroidal reaction product with an acylating agent effective to replace the hydroxy group by a said lower acyloxy group.